

Bibliography

- [1] S. Aaronson. *Quantum computing since Democritus*. Cambridge University Press, Cambridge, 2013.
- [2] E. Abdallah, A. Hamza, and P. Bhattacharya. Mpeg video watermarking using tensor singular value decomposition. In *Image Analysis and Recognition*, volume 4633 of *Lecture Notes in Computer Science*, pages 772–783. Springer Berlin / Heidelberg, 2007. 10.1007/978-3-540-74260-9_69.
- [3] P.-A. Absil, R. Mahony, and B. Andrews. Convergence of the iterates of descent methods for analytic cost functions. *SIAM J. Optim.*, 16(2):531–547, 2005.
- [4] E. Acar, C. Aykut-Bingol, H. Bingol, R. Bro, and B. Yener. Multiway analysis of epilepsy tensors. *Bioinformatics*, 23(13):i10–i18, July 2007.
- [5] E. Acar, C. A. Bingol, H. Bingol, R. Bro, and B. Yener. Seizure recognition on epilepsy feature tensor. In *EMBS 2007: Proceedings of the 29th Annual International Conference on Engineering in Medicine and Biology Society, IEEE*, pages 4273–4276.
- [6] E. Acar, S. A. Çamtepe, and B. Yener. *Collective Sampling and Analysis of High Order Tensors for Chatroom Communications*, pages 213–224. ISI'06. Springer Berlin Heidelberg, Berlin, Heidelberg, 2006.
- [7] E. Acar, S. A. Çamtepe, M. S. Krishnamoorthy, and B. Yener. Modeling and multiway analysis of chatroom tensors. In *Proceedings of the 2005 IEEE international conference on Intelligence and Security Informatics, ISI'05*, pages 256–268, Berlin, Heidelberg, 2005. Springer-Verlag.
- [8] E. Acar, D. M. Dunlavy, and T. G. Kolda. A scalable optimization approach for fitting canonical tensor decompositions. *Journal of Chemometrics*, 25(2):67–86, February 2011.
- [9] E. Acar and B. Yener. Unsupervised multiway data analysis: A literature survey. *IEEE Trans. on Knowl. and Data Eng.*, 21(1):6–20, Jan. 2009.
- [10] S. N. Afriat. Theory of maxima and the method of Lagrange. *SIAM J. Appl. Math.*, 20:343–357, 1971.
- [11] L. Albera, A. Ferréol, P. Comon, and P. Chevalier. Blind identification of overcomplete mixtures of sources (BIOME). *Linear Algebra Appl.*, 391:3–30, 2004.

- [12] O. Alter, P. O. Brown, and D. Botstein. Singular value decomposition for genome-wide expression data processing and modeling. *Proceedings of The National Academy of Sciences*, 97:10101–10106, 2000.
- [13] A. H. Andersen and W. S. Rayens. Structure-seeking multilinear methods for the analysis of fmri data. *NeuroImage*, 22(2):728 – 739, 2004.
- [14] C. M. Andersen and R. Bro. Practical aspects of parafac modeling of fluorescence excitation-emission data. *Journal of Chemometrics*, 17(4):200–215, 2003.
- [15] C. A. Andersson and R. Bro. Improving the speed of multi-way algorithms: - part I. Tucker3. *Chemometrics and Intelligent Laboratory Systems*, 42(1):93–103, 1998.
- [16] C. A. Andersson and R. Bro. The n-way toolbox for matlab. *Chemometrics and Intelligent Laboratory Systems*, 52(1):1 – 4, 2000.
- [17] C. A. Andersson and R. Henrion. A general algorithm for obtaining simple structure of core arrays in n-way pca with application to fluorometric data. *Computational Statistics and Data Analysis*, 31(3):255 – 278, 1999.
- [18] P. Aniello, J. Clemente-Gallardo, G. Marmo, and G. F. Volkert. Classical tensors and quantum entanglement II: mixed states. *Int. J. Geom. Methods Mod. Phys.*, 8(4):853–883, 2011.
- [19] C. J. Appellof and E. R. Davidson. Strategies for analyzing data from video fluorometric monitoring of liquid chromatographic effluents. *Analytical Chemistry*, 53(13):2053–2056, 1981.
- [20] V. I. Arnold. *Real algebraic geometry*, volume 66 of *Unitext*. Springer, Heidelberg, 2013. Translated from the 2009 Russian original by Gerald G. Gould and David Kramer.
- [21] H. Attouch and J. Bolte. On the convergence of the proximal algorithm for nonsmooth functions involving analytic features. *Math. Program.*, 116(1-2, Ser. B):5–16, 2009.
- [22] H. Attouch, J. Bolte, and B. F. Svaiter. Convergence of descent methods for semi-algebraic and tame problems: proximal algorithms, forward-backward splitting, and regularized Gauss-Seidel methods. *Math. Program.*, 137(1-2, Ser. A):91–129, 2013.
- [23] J. E. Avron, G. Bisker, and O. Kenneth. Visualizing two qubits. *J. Math. Phys.*, 48(10):102107, 14, 2007.
- [24] B. Bader, R. Harshman, and T. Kolda. Temporal analysis of semantic graphs using asalsan. In *Data Mining, 2007. ICDM 2007. Seventh IEEE International Conference on*, pages 33–42, oct. 2007.
- [25] B. W. Bader, M. W. Berry, and M. Browne. Discussion tracking in enron email using parafac. In M. W. Berry and M. Castellanos, editors, *Survey of Text Mining II*, pages 147–163. Springer London, 2008. 10.1007/978-1-84800-046-9_8.
- [26] B. W. Bader and T. G. Kolda. Algorithm 862: MATLAB tensor classes for fast algorithm prototyping. *ACM Trans. Math. Software*, 32(4):635–653, 2006.

- [27] B. W. Bader and T. G. Kolda. Efficient matlab computations with sparse and factored tensors. *SIAM J. Sci. Comput.*, 30(1):205–231, Dec. 2007.
- [28] B. W. Bader and T. G. Kolda. Efficient MATLAB computations with sparse and factored tensors. *SIAM J. Sci. Comput.*, 30(1):205–231, 2007/08.
- [29] S. Banach. Über homogene polynome in (L^2) . *Studia Mathematica*, 7(1):36–44, 1938.
- [30] S. Basu, R. Pollack, and M.-F. Roy. *Algorithms in real algebraic geometry*, volume 10 of *Algorithms and Computation in Mathematics*. Springer-Verlag, Berlin, second edition, 2006.
- [31] K. Batselier, H. Liu, and N. Wong. A constructive algorithm for decomposing a tensor into a finite sum of orthonormal rank-1 terms. *SIAM J. Matrix Anal. Appl.*, 36(3):1315–1337, 2015.
- [32] K. Batselier and N. Wong. Symmetric tensor decomposition by an iterative eigendecomposition algorithm. *J. Comput. Appl. Math.*, 308:69–82, 2016.
- [33] C. Bauckhage. Robust tensor classifiers for color object recognition. In M. Kamel and A. Campilho, editors, *Image Analysis and Recognition*, volume 4633 of *Lecture Notes in Computer Science*, pages 352–363. Springer Berlin / Heidelberg, 2007. 10.1007/978-3-540-74260-9_32.
- [34] C. Beckmann and S. Smith. Tensorial extensions of independent component analysis for multisubject fmri analysis. *NeuroImage*, 25(1):294 – 311, 2005.
- [35] J. S. Bell. On the einstein podolsky rosen paradox. *Physics Physique Fizika*, 1:195–200, Nov 1964.
- [36] C. H. Bennett, G. Brassard, C. Crépeau, R. Jozsa, A. Peres, and W. K. Wootters. Teleporting an unknown quantum state via dual classical and einstein-podolsky-rosen channels. *Phys. Rev. Lett.*, 70:1895–1899, Mar 1993.
- [37] C. H. Bennett and S. J. Wiesner. Communication via one- and two-particle operators on einstein-podolsky-rosen states. *Phys. Rev. Lett.*, 69:2881–2884, Nov 1992.
- [38] J. M. T. Berge and H. A. Kiers. Simplicity of core arrays in three-way principal component analysis and the typical rank of $p \times q \times 2$ arrays. *Linear Algebra Appl.*, 294(1&C3):169 – 179, 1999.
- [39] M. W. Berry, S. T. Dumais, and G. W. O’Brien. Using Linear Algebra for Intelligent Information Retrieval. Technical Report UT-CS-94-270, 1995.
- [40] R. A. Bertlmann and P. Krammer. Entanglement witnesses and geometry of entanglement of two-qutrit states. *Ann. Physics*, 324(7):1388–1407, 2009.
- [41] G. Beylkin and M. J. Mohlenkamp. Numerical operator calculus in higher dimensions. *Proceedings of the National Academy of Sciences*, 99(16):10246–10251, 2002.
- [42] G. Beylkin and M. J. Mohlenkamp. Algorithms for numerical analysis in high dimensions. *SIAM J. Sci. Comput.*, 26(6):2133–2159, June 2005.

- [43] J. C. Bezdek and R. J. Hathaway. Some notes on alternating optimization. In N. R. Pal and M. Sugeno, editors, *Advances in Soft Computing — AFSS 2002: 2002 AFSS International Conference on Fuzzy Systems Calcutta, India.*, pages 288–300. Springer, Berlin, Heidelberg, 2002.
- [44] J. C. Bezdek and R. J. Hathaway. Convergence of alternating optimization. *Neural Parallel Sci. Comput.*, 11(4):351–368, 2003.
- [45] D. Bini, M. Capovani, F. Romani, and G. Lotti. $O(n^{2.7799})$ Complexity for $n \times n$ Approximate Matrix Multiplication. *Information Processing Letters*, 8:234–235, 1979.
- [46] M. Bismarck. Multiway data analysis: Coppi, r., bolasco, s. (eds.) (1989): Amsterdam, new york, oxford, tokyo: North-holland. 552 pp isbn 0-444-87410-0. *Computational Statistics and Data Analysis*, 11(3):356–357, May 1991.
- [47] M. Bläser. On the complexity of the multiplication of matrices of small formats. *Journal of Complexity*, 19(1):43 – 60, 2003.
- [48] M. Bolten, K. Kahl, and S. Sokolović. Multigrid methods for tensor structured Markov chains with low rank approximation. *SIAM J. Sci. Comput.*, 38(2):A649–A667, 2016.
- [49] S. Boyd and L. Xiao. Least-squares covariance matrix adjustment. *SIAM J. Matrix Anal. Appl.*, 27(2):532–546, 2005.
- [50] J. Brachat, P. Comon, B. Mourrain, and E. Tsigaridas. Symmetric tensor decomposition. *Linear Algebra Appl.*, 433(11-12):1851–1872, 2010.
- [51] D. H. Brandwood. A complex gradient operator and its application in adaptive array theory. *Proc. IEE-H*, 130(1):11–16, 1983.
- [52] G. E. Bredon. *Introduction to compact transformation groups*. Academic Press, New York-London, 1972. Pure and Applied Mathematics, Vol. 46.
- [53] R. Bro. Parafac. tutorial and applications. *Chemometrics and Intelligent Laboratory Systems*, 38(2):149 – 171, 1997.
- [54] R. Bro. *Multi-way analysis in the food industry. Models, algorithms, and applications*. PhD thesis, University of Amsterdam, Amsterdam, The Netherlands, 1998.
- [55] R. Bro. Review on multiway analysis in chemistry—2002-2005. *Critical Reviews in Analytical Chemistry*, 36(3-4):279–293, 2006.
- [56] R. Bro, C. A. Andersson, and H. A. L. Kiers. Parafac2^oTM part ii. modeling chromatographic data with retention time shifts. *Journal of Chemometrics*, 13(3-4):295–309, 1999.
- [57] R. Bro and S. De Jong. A fast non-negativity-constrained least squares algorithm. *Journal of Chemometrics*, 11(5):393–401, 1997.
- [58] R. Bro, R. Harshman, N. Sidiropoulos, and M. Lundy. Modeling multi-way data with linearly dependent loadings. *Journal of Chemometrics*, 23(7-8):324–340, 2009.

- [59] R. Bro and H. A. L. Kiers. A new efficient method for determining the number of components in parafac models. *Journal of Chemometrics*, 17(5):274–286, 2003.
- [60] R. Bro, N. D. Sidiropoulos, and A. K. Smilde. Maximum likelihood fitting using ordinary least squares algorithms. *Journal of Chemometrics*, 16(8-10):387–400, 2002.
- [61] R. Bro and A. K. Smilde. Centering and scaling in component analysis. *Journal of Chemometrics*, 17(1):16–33, 2003.
- [62] D. Bruß. Characterizing entanglement. *Journal of Mathematical Physics*, 43(9):4237–4251, 2002.
- [63] N. Bshouty. Maximal rank of $m \times n \times (mn - k)$ tensors. *SIAM Journal on Computing*, 19(3):467–471, 1990.
- [64] A. Bunse-Gerstner, R. Byers, V. Mehrmann, and N. K. Nichols. Numerical computation of an analytic singular value decomposition of a matrix valued function. *Numer. Math.*, 60:1–40, 1991.
- [65] J. A. Cadzow. Signal enhancement: A composite property mapping algorithm. *IEEE Trans. on Acoust., Speech, Signal Processing*, 36:39–62, 1988.
- [66] M. E. Carrington, R. Kobes, G. Kunstatter, D. Ostapchuk, and G. Passante. Geometric measures of entanglement and the Schmidt decomposition. *J. Phys. A*, 43(31):315302, 15, 2010.
- [67] J. Carroll and J.-J. Chang. Analysis of individual differences in multidimensional scaling via an n-way generalization of Eckart-Young decomposition. *Psychometrika*, 35:283–319, 1970.
- [68] R. Cattell. Parallel proportional profiles and other principles for determining the choice of factors by rotation. *Psychometrika*, 9(4):267–283, 1944.
- [69] R. B. Cattell. The three basic factor-analytic research designs^{oTM}their interrelations and derivatives. *Psychological Bulletin*, 49(5):499 – 520, 1952.
- [70] S. Celar, Z. Stojkic, Z. Seremet, Z. Marusic, and D. Zelenika. Classification of test documents based on handwritten student id’s characteristics. *Procedia Engineering*, 100(Supplement C):782 – 790, 2015. 25th DAAAM International Symposium on Intelligent Manufacturing and Automation, 2014.
- [71] N. J. Cerf, C. Adami, and R. M. Gingrich. Reduction criterion for separability. *Phys. Rev. A* (3), 60(2):898–909, 1999.
- [72] E. Ceulemans and H. A. L. Kiers. Selecting among three-mode principal component models of different types and complexities: A numerical convex hull based method. *British Journal of Mathematical and Statistical Psychology*, 59(1):133–150, 2006.
- [73] J. Chang, W. Sun, and Y. Chen. A modified newton’s method for best rank-one approximation to tensors. *Applied Mathematics and Computation*, 216(6):1859 – 1867, 2010. Workshop on Iterative Methods and Preconditioning Techniques.

- [74] K. Chang, L. Qi, and T. Zhang. A survey on the spectral theory of nonnegative tensors. *Numer. Linear Algebra Appl.*, 20(6):891–912, 2013.
- [75] B. Chen, Z. Li, and S. Zhang. On optimal low rank Tucker approximation for tensors: the case for an adjustable core size. *J. Global Optim.*, 62(4):811–832, 2015.
- [76] B. Chen, A. P. Petropulu, and L. De Lathauwer. Blind identification of convolutive mimo systems with 3 sources and 2 sensors. *EURASIP J. Appl. Signal Process.*, 2002(1):487–496, Jan. 2002.
- [77] H. Chen. Constraints on the mixing of bipartite states. *J. Math. Phys.*, 46(12):122101, 7, 2005.
- [78] H. Chen. Quantum entanglement and geometry of determinantal varieties. *J. Math. Phys.*, 47(5):052101, 19, 2006.
- [79] J. Chen and Y. Saad. On the tensor SVD and the optimal low rank orthogonal approximation of tensors. *SIAM J. Matrix Anal. Appl.*, 30(4):1709–1734, 2008/09.
- [80] K. Chen and L.-A. Wu. A matrix realignment method for recognizing entanglement. *Quantum Inf. Comput.*, 3(3):193–202, 2003.
- [81] L. Chen, M. Aulbach, and M. Hajdušek. Comparison of different definitions of the geometric measure of entanglement. *Phys. Rev. A*, 89:042305, Apr 2014.
- [82] L. Cheng, Y.-C. Wu, and H. V. Poor. Probabilistic tensor canonical polyadic decomposition with orthogonal factors. *IEEE Trans. Signal Process.*, 65(3):663–676, 2017.
- [83] P. A. Chew, B. W. Bader, T. G. Kolda, and A. Abdelali. Cross-language information retrieval using parafac2. In *Proceedings of the 13th ACM SIGKDD international conference on Knowledge discovery and data mining*, KDD '07, pages 143–152, New York, NY, USA, 2007. ACM.
- [84] R. Chill. On the Łojasiewicz-Simon gradient inequality. *J. Funct. Anal.*, 201(2):572–601, 2003.
- [85] E. Chitambar, C. A. Miller, and Y. Shi. Matrix pencils and entanglement classification. *J. Math. Phys.*, 51(7):072205, 16, 2010.
- [86] M. T. Chu. On the continuous realization of iterative processes. *SIAM Review*, 30(3):375–387, 1988.
- [87] M. T. Chu and K. R. Driessel. The projected gradient method for least squares matrix approximations with spectral constraints. *SIAM J. Numer. Anal.*, 27(4):1050–1060, 1990.
- [88] M. T. Chu, R. E. Funderlic, and G. H. Golub. A rank-one reduction formula and its applications to matrix factorizations. *SIAM Rev.*, 37(4):512–530, 1995.
- [89] M. T. Chu, R. E. Funderlic, and R. J. Plemmons. Structured low rank approximation. *Linear Algebra Appl.*, 366:157–172, 2003. Special issue on structured matrices: analysis, algorithms and applications (Cortona, 2000).

- [90] M. T. Chu and M. H. Lin. Rank-1 approximation for bipartite entangled systems. preprint, North Carolina State University, 2019.
- [91] M. T. Chu and M. M. Lin. Convergence analysis of alternating direction methods: A general framework and its applications to tensor approximations. submitted, North Carolina State University, 2020.
- [92] M. T. Chu, M. M. Lin, and L. Wang. A study of singular spectrum analysis with global optimization techniques. *J. Global Optim.*, 60(3):551–574, 2014.
- [93] M. T. Chu and N. T. Trendafilov. The orthogonally constrained regression revisited. *J. Comput. Graph. Statist.*, 10(4):746–771, 2001.
- [94] M. T. Chu and J. L. Watterson. On a multivariate eigenvalue problem. I. Algebraic theory and a power method. *SIAM J. Sci. Comput.*, 14(5):1089–1106, 1993.
- [95] A. Cichocki, R. Zdunek, S. Choi, R. Plemmons, and S. Amari. Non-negative tensor factorization using alpha and beta divergences. In *Acoustics, Speech and Signal Processing, 2007. ICASSP 2007. IEEE International Conference on*, volume 3, pages III–1393 –III–1396, april 2007.
- [96] P. Comon. Independent component analysis, a new concept? *Signal Process.*, 36(3):287–314, Apr. 1994.
- [97] P. Comon. Tensor decompositions: state of the art and applications. In *Mathematics in signal processing, V (Coventry, 2000)*, volume 71 of *Inst. Math. Appl. Conf. Ser. New Ser.*, pages 1–24. Oxford Univ. Press, Oxford, Oxford, 2002.
- [98] P. Comon, G. Golub, L.-H. Lim, and B. Mourrain. Symmetric tensors and symmetric tensor rank. *SIAM J. Matrix Anal. Appl.*, 30(3):1254–1279, Sept. 2008.
- [99] P. Comon, X. Luciani, and A. L. F. de Almeida. Tensor decompositions, alternating least squares and other tales. *J. Chemometrics*, 23:393–405, 2009.
- [100] P. Comon and B. Mourrain. Decomposition of quantics in sums of powers of linear forms. *Signal Process.*, 53(2-3):93–107, Sept. 1996.
- [101] P. Comon, J. M. F. ten Berge, L. De Lathauwer, and J. Castaing. Generic and typical ranks of multi-way arrays. *Linear Algebra Appl.*, 430(11-12):2997–3007, 2009.
- [102] F. Cong, Q.-H. Lin, L.-D. Kuang, X.-F. Gong, P. Astikainen, and T. Ristaniemi. Tensor decomposition of eeg signals: A brief review. *Journal of Neuroscience Methods*, 248(Supplement C):59 – 69, 2015.
- [103] A. P. da Silva, P. Comon, and A. L. F. de Almeida. A finite algorithm to compute rank-1 tensor approximations. *IEEE Signal Processing Letters*, 23(7):959–963, July 2016.
- [104] M. Dana and K. D. Ikramov. On the codimension of the variety of symmetric matrices with multiple eigenvalues. *Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI)*, 323(Chisl. Metody i Vopr. Organ. Vychisl. 18):34–46, 224, 2005.

- [105] W. F. de la Vega, M. Karpinski, R. Kannan, and S. Vempala. Tensor decomposition and approximation schemes for constraint satisfaction problems. In *Proceedings of the thirty-seventh annual ACM symposium on Theory of computing*, STOC '05, pages 747–754, New York, NY, USA, 2005. ACM.
- [106] L. De Lathauwer. A link between the canonical decomposition in multilinear algebra and simultaneous matrix diagonalization. *SIAM J. Matrix Anal. Appl.*, 28(3):642–666, Aug. 2006.
- [107] L. De Lathauwer. Decompositions of a higher-order tensor in block terms – part ii: Definitions and uniqueness. *SIAM J. Matrix Anal. Appl.*, 30(3):1033–1066, Sept. 2008.
- [108] L. De Lathauwer. Decompositions of a higher-order tensor in block terms –part i: Lemmas for partitioned matrices. *SIAM J. Matrix Anal. Appl.*, 30(3):1022–1032, Sept. 2008.
- [109] L. De Lathauwer and J. Castaing. Tensor-based techniques for the blind separation of ds-cdma signals. *Signal Process.*, 87(2):322–336, Feb. 2007.
- [110] L. De Lathauwer and J. Castaing. Blind identification of underdetermined mixtures by simultaneous matrix diagonalization. *IEEE Trans. Signal Process.*, 56(3):1096–1105, 2008.
- [111] L. De Lathauwer, J. Castaing, and J.-F. Cardoso. Fourth-order cumulant-based blind identification of underdetermined mixtures. *IEEE Trans. Signal Process.*, 55(6, part 2):2965–2973, 2007.
- [112] L. De Lathauwer, P. Comon, and N. Mastronardi. Special issue on tensor decompositions and applications. *SIAM J. Matrix Anal. Appl.*, 30(3):vii, 2008.
- [113] L. De Lathauwer and A. de Baynast. Blind deconvolution of DS-CDMA signals by means of decomposition in rank- $(1, L, L)$ terms. *IEEE Trans. Signal Process.*, 56(4):1562–1571, 2008.
- [114] L. De Lathauwer, B. De Moor, and J. Vandewalle. A multilinear singular value decomposition. *SIAM J. Matrix Anal. Appl.*, 21(4):1253–1278 (electronic), 2000.
- [115] L. De Lathauwer, B. De Moor, and J. Vandewalle. On the best rank-1 and rank- (r_1, r_2, \dots, r_n) approximation of higher-order tensors. *SIAM journal on Matrix Analysis and Applications*, 21(4):1324–1342, 2000.
- [116] L. De Lathauwer, B. De Moor, and J. Vandewalle. On the best rank-1 and rank- (R_1, R_2, \dots, R_N) approximation of higher-order tensors. *SIAM J. Matrix Anal. Appl.*, 21(4):1324–1342 (electronic), 2000.
- [117] L. De Lathauwer, B. de Moor, and J. Vandewalle. Independent component analysis and (simultaneous) third-order tensor diagonalization. *Signal Processing, IEEE Transactions on*, 49(10):2262–2271, oct 2001.
- [118] L. De Lathauwer, B. De Moor, and J. Vandewalle. Computation of the canonical decomposition by means of a simultaneous generalized Schur decomposition. *SIAM J. Matrix Anal. Appl.*, 26(2):295–327 (electronic), 2004/05.

- [119] L. De Lathauwer and D. Nion. Decompositions of a higher-order tensor in block terms — part iii: Alternating least squares algorithms. *SIAM J. Matrix Anal. Appl.*, 30(3):1067–1083, Sept. 2008.
- [120] L. De Lathauwer and J. Vandewalle. Dimensionality reduction in higher-order signal processing and rank- (R_1, R_2, \dots, R_N) reduction in multilinear algebra. *Linear Algebra Appl.*, 391:31–55, 2004.
- [121] V. de Silva and L.-H. Lim. Tensor rank and the ill-posedness of the best low-rank approximation problem. *SIAM J. Matrix Anal. Appl.*, 30(3):1084–1127, 2008.
- [122] H. De Sterck and K. Miller. An adaptive algebraic multigrid algorithm for low-rank canonical tensor decomposition. *SIAM J. Sci. Comput.*, 35(1):B1–B24, 2013.
- [123] M. De Vos, L. De Lathauwer, B. Vanrumste, S. Van Huffel, and W. Van Paesschen. Canonical decomposition of ictal scalp EEG and accurate source localisation: principles and simulation study. *Intell. Neuroscience*, 2007:11:1–11:8, Apr. 2007.
- [124] S. Deerwester, S. T. Dumais, G. W. Furnas, T. K. Landauer, and R. Harshman. Indexing by latent semantic analysis. *Journal of the American Society for Information Science*, 41(6):391–407, 1990.
- [125] R. S. Dembo, S. C. Eisenstat, and T. Steihaug. Inexact Newton methods. *SIAM J. Numer. Anal.*, 19(2):400–408, 1982.
- [126] L. u. Derkacz, M. Gwóźdź, and L. Jakóbczyk. Entanglement beyond tensor product structure: algebraic aspects of quantum non-separability. *J. Phys. A*, 45(2):025302, 11, 2012.
- [127] H. Derksen and G. Kemper. *Computational invariant theory*, volume 130 of *Encyclopaedia of Mathematical Sciences*. Springer, Heidelberg, enlarged edition, 2015. Invariant Theory and Algebraic Transformation Groups, VIII.
- [128] I. Domanov and L. De Lathauwer. On the uniqueness of the canonical polyadic decomposition of third-order tensors—Part I: Basic results and uniqueness of one factor matrix. *SIAM J. Matrix Anal. Appl.*, 34(3):855–875, 2013.
- [129] B. Dong, M. M. Lin, and M. T. Chu. Nonnegative rank factorization—a heuristic approach via rank reduction. *Numer. Algorithms*, 65(2):251–274, 2014.
- [130] J. Douglas Carroll, S. Pruzansky, and J. Kruskal. Candelinec: A general approach to multi-dimensional analysis of many-way arrays with linear constraints on parameters. *Psychometrika*, 45:3–24, 1980. 10.1007/BF02293596.
- [131] P. Drineas and M. W. Mahoney. A randomized algorithm for a tensor-based generalization of the singular value decomposition. *Linear Algebra Appl.*, 420:553–571, 2007.
- [132] R. Duan, Y. Feng, Y. Xin, and M. Ying. Distinguishability of quantum states by separable operations. *IEEE Trans. Inform. Theory*, 55(3):1320–1330, 2009.

- [133] S. T. Dumais, G. W. Furnas, T. K. Landauer, S. Deerwester, and R. Harshman. Using latent semantic analysis to improve access to textual information. In *Proceedings of the SIGCHI conference on Human factors in computing systems*, CHI '88, pages 281–285, New York, NY, USA, 1988. ACM.
- [134] C. Eckart and G. Young. The approximation of one matrix by another of lower rank. *Psychometrika*, 1(3):211–218, 1936.
- [135] A. Einstein, B. Podolsky, and N. Rosen. Can quantum-mechanical description of physical reality be considered complete? *Phys. Rev.*, 47:777–780, May 1935.
- [136] S. C. Eisenstat and H. F. Walker. Globally convergent inexact Newton methods. *SIAM J. Optim.*, 4(2):393–422, 1994.
- [137] J. Eisert, K. Audenaert, and M. B. Plenio. Remarks on entanglement measures and non-local state distinguishability. *J. Phys. A*, 36(20):5605–5615, 2003.
- [138] A. Ekert and P. L. Knight. Entangled quantum systems and the schmidt decomposition. *American Journal of Physics*, 63(5):415–423, 1995.
- [139] A. K. Ekert. Quantum cryptography based on bell’s theorem. *Phys. Rev. Lett.*, 67:661–663, Aug 1991.
- [140] L. Eldén and B. Savas. A Newton-Grassmann method for computing the best multilinear rank- (r_1, r_2, r_3) approximation of a tensor. *SIAM J. Matrix Anal. Appl.*, 31(2):248–271, 2009.
- [141] M. Espig and A. Khachatryan. Convergence of Alternating Least Squares Optimisation for Rank-One Approximation to High Order Tensors. *arXiv e-prints*, page arXiv:1503.05431, Mar 2015.
- [142] N. K. M. Faber, R. Bro, and P. K. Hopke. Recent developments in candecomp/parafac algorithms: a critical review. *Chemometrics and Intelligent Laboratory Systems*, 65(1):119 – 137, 2003.
- [143] A. S. Field and D. Graupe. Topographic component (parallel factor) analysis of multichannel evoked potentials: Practical issues in trilinear spatiotemporal decomposition. *Brain Topography*, 3:407–423, 1991. 10.1007/BF01129000.
- [144] S. Friedland. Best rank one approximation of real symmetric tensors can be chosen symmetric. *Front. Math. China*, 8(1):19–40, 2013.
- [145] S. Friedland and G. Gour. An explicit expression for the relative entropy of entanglement in all dimensions. *J. Math. Phys.*, 52(5):052201, 13, 2011.
- [146] S. Friedland, V. Mehrmann, R. Pajarola, and S. K. Suter. On best rank one approximation of tensors. *Numer. Linear Algebra Appl.*, 20(6):942–955, 2013.
- [147] M. P. Friedlander and K. Hatz. Computing nonnegative tensor factorizations. *Computational Optimization and Applications*, 23(4):631–647, March 2008.

- [148] R. Furukawa, H. Kawasaki, K. Ikeuchi, and M. Sakauchi. Appearance based object modeling using texture database: acquisition, compression and rendering. In *Proceedings of the 13th Eurographics workshop on Rendering, EGRW '02*, pages 257–266, Aire-la-Ville, Switzerland, Switzerland, 2002. Eurographics Association.
- [149] C. B. García and T.-Y. Li. On the number of solutions to polynomial systems of equations. *SIAM J. Numer. Anal.*, 17(4):540–546, 1980.
- [150] R. Garcia and A. Lumsdaine. Multiarray: a c++ library for generic programming with arrays. *Software: Practice and Experience*, 35(2):159–188, 2005.
- [151] S. Gharibian. Strong np-hardness of the quantum separability problem. *Quantum Info. Comput.*, 10(3):343–360, Mar. 2010.
- [152] J. Gillard and A. Zhigljavsky. Application of structured low-rank approximation methods for imputing missing values in the time series. *Stat. Interface*, 8(3):321–330, 2015.
- [153] G. H. Golub and C. F. Van Loan. *Matrix computations*. Johns Hopkins Studies in the Mathematical Sciences. Johns Hopkins University Press, Baltimore, MD, fourth edition, 2013.
- [154] G. H. Golub and C. F. van Van Loan. *Matrix Computations (Johns Hopkins Studies in Mathematical Sciences)(3rd Edition)*. The Johns Hopkins University Press, 3rd edition, Oct. 1996.
- [155] S. A. Goreinov, I. V. Oseledets, and D. V. Savostyanov. Wedderburn rank reduction and Krylov subspace method for tensor approximation. Part 1: Tucker case. *SIAM J. Sci. Comput.*, 34(1):A1–A27, 2012.
- [156] L. Grasedyck. Existence and computation of low kronecker-rank approximations for large linear systems of tensor product structure. *Computing*, 72:247–265, 2004. 10.1007/s00607-003-0037-z.
- [157] L. Grasedyck. Hierarchical singular value decomposition of tensors. *SIAM Journal on Matrix Analysis and Applications*, 31(4):2029–2054, 2010.
- [158] L. Grippo and M. Sciandrone. On the convergence of the block nonlinear Gauss-Seidel method under convex constraints. *Oper. Res. Lett.*, 26(3):127–136, 2000.
- [159] W. Gropp and J. J. Moré. Optimization environments and the neos server. In M. D. Buhman and A. Iserles, editors, *Approximation Theory and Optimization*, pages 167 – 182. Cambridge University Press, 1997.
- [160] Y. Guan, M. T. Chu, and D. Chu. Convergence analysis of an SVD-based algorithm for the best rank-1 tensor approximation. *Linear Algebra Appl.*, 555:53–69, 2018.
- [161] Y. Guan, M. T. Chu, and D. Chu. SVD-based algorithms for the best rank-1 approximation of a symmetric tensor. *SIAM J. Matrix Anal. Appl.*, 39(3):1095–1115, 2018.
- [162] O. Gühne and G. Tóth. Entanglement detection. *Phys. Rep.*, 474(1-6):1–75, 2009.

- [163] V. Guillemin and A. Pollack. *Differential topology*. AMS Chelsea Publishing, Providence, RI, 2010. Reprint of the 1974 original.
- [164] L. Gurvits. Classical deterministic complexity of edmonds' problem and quantum entanglement. In *Proceedings of the Thirty-fifth Annual ACM Symposium on Theory of Computing, STOC '03*, pages 10–19, New York, NY, USA, 2003. ACM.
- [165] K.-C. Ha and S.-H. Kye. The structural physical approximations and optimal entanglement witnesses. *J. Math. Phys.*, 53(10):102204, 7, 2012.
- [166] W. Hackbusch. *Tensor spaces and numerical tensor calculus*, volume 42 of *Springer Series in Computational Mathematics*. Springer, Heidelberg, 2012.
- [167] W. Hackbusch. New estimates for the recursive low-rank truncation of block-structured matrices. *Numer. Math.*, 132(2):303–328, 2016.
- [168] W. Hackbusch and B. N. Khoromskij. Tensor-product approximation to operators and functions in high dimensions. *J. Complexity*, 23(4-6):697–714, 2007.
- [169] W. Hackbusch, B. N. Khoromskij, and E. E. Tyrtyshnikov. Hierarchical Kronecker Tensor-Product Approximations. 2003.
- [170] W. Hackbusch, B. N. Khoromskij, and E. E. Tyrtyshnikov. Hierarchical Kronecker tensor-product approximations. *J. Numer. Math.*, 13(2):119–156, 2005.
- [171] K. H. Han and S.-H. Kye. Various notions of positivity for bi-linear maps and applications to tri-partite entanglement. *J. Math. Phys.*, 57(1):015205, 22, 2016.
- [172] R. Harshman. Foundations of the parafac procedure: Models and conditions for an “explanatory” multi-modal factor analysis. *UCLA Working Papers in Phonetics*, 16, 1970.
- [173] R. A. Harshman. Determination and proof of minimum uniqueness conditions for PARAFAC1. *UCLA Working Papers in Phonetics*, 22, 1972.
- [174] R. A. Harshman. PARAFAC2: Mathematical and technical notes. *UCLA Working Papers in Phonetics*, 22:30–44, 1972b.
- [175] R. A. Harshman. An index formalism that generalizes the capabilities of matrix notation and algebra to n-way arrays. *Journal of Chemometrics*, 15(9):689–714, 2001.
- [176] R. A. Harshman, P. E. Green, Y. Wind, and M. E. Lundy. A model for the analysis of asymmetric data in marketing research. *Marketing Science*, 1(2):205–242, 1982.
- [177] R. A. Harshman and S. Hong. ”Stretch” vs ”slice” methods for representing three-way structure via matrix notation. *Journal of Chemometrics*, 16(4):198–205, 2002.
- [178] R. A. Harshman, S. Hong, and M. E. Lundy. Shifted factor analysis – part i: Models and properties, 2003.
- [179] R. A. Harshman and M. E. Lundy. Data preprocessing and the extended PARAFAC model. In H. G. Law, C. W. Snyder Jr, J. A. Hattie, and R. P. McDonald, editors, *Research methods for multimode data analysis.*, pages 216–284. Praeger, New York, 1984b.

- [180] R. A. Harshman, Margaret, and E. Lundy. Uniqueness proof for a family of models sharing features of tucker's three-mode factor analysis and parafac/candecomp. *Psychometrika*, 1996.
- [181] R. Hartshorne. *Algebraic geometry*. Springer-Verlag, New York, 1977. Graduate Texts in Mathematics, No. 52.
- [182] J. Håstad. Tensor rank is NP-complete. *J. Algorithms*, 11(4):644–654, Dec. 1990.
- [183] R. J. Hathaway, Y. Hu, and J. C. Bezdek. Local convergence of tri-level alternating optimization. *Neural Parallel Sci. Comput.*, 9(1):19–28, 2001.
- [184] M. Hayashi. *Quantum information theory*. Graduate Texts in Physics. Springer-Verlag, Berlin, second edition, 2017. Mathematical foundation.
- [185] M. Hayashi, D. Markham, M. Muraio, M. Owari, and S. Virmani. The geometric measure of entanglement for a symmetric pure state with non-negative amplitudes. *J. Math. Phys.*, 50(12):122104, 6, 2009.
- [186] T. Hazan, S. Polak, and A. Shashua. Sparse image coding using a 3d non-negative tensor factorization. In *Proceedings of the Tenth IEEE International Conference on Computer Vision (ICCV'05) Volume 1 - Volume 01*, ICCV '05, pages 50–57, Washington, DC, USA, 2005. IEEE Computer Society.
- [187] M. Hein and T. Bühler. An inverse power method for nonlinear eigenproblems with applications in 1-spectral clustering and sparse PCA. *CoRR*, abs/1012.0774, 2010.
- [188] R. Henrion. Body diagonalization of core matrices in three-way principal components analysis: Theoretical bounds and simulation. *Journal of Chemometrics*, 7(6):477–494, 1993.
- [189] R. Henrion. N-way principal component analysis theory, algorithms and applications. *Chemometrics and Intelligent Laboratory Systems*, 25(1):1 – 23, 1994.
- [190] F. Hiai and D. Petz.
- [191] N. J. Higham. Computing the polar decomposition—with applications. *SIAM J. Sci. Statist. Comput.*, 7(4):1160–1174, 1986.
- [192] N. J. Higham. Computing a nearest symmetric positive semidefinite matrix. *Linear Algebra Appl.*, 103:103–118, 1988.
- [193] N. J. Higham. Computing the nearest correlation matrix—a problem from finance. *IMA J. Numer. Anal.*, 22(3):329–343, 2002.
- [194] C. J. Hillar and L.-H. Lim. Most tensor problems are NP-hard. *J. ACM*, 60(6):Art. 45, 39, 2013.
- [195] J. J. Hilling and A. Sudbery. The geometric measure of multipartite entanglement and the singular values of a hypermatrix. *J. Math. Phys.*, 51(7):072102, 15, 2010.

- [196] T. Hiroshima. Majorization criterion for distillability of a bipartite quantum state. *Phys. Rev. Lett.*, 91:057902, Aug 2003.
- [197] F. Hitchcock. *The Expression of a Tensor Or a Polyadic as a Sum of Products*. Contributions from the Department of Mathematics. sn., 1927.
- [198] M. Horodecki, P. Horodecki, and R. Horodecki. Separability of mixed states: necessary and sufficient conditions. *Physics Letters A*, 223(1):1 – 8, 1996.
- [199] M. Horodecki, P. Horodecki, and R. Horodecki. *Mixed-State Entanglement and Quantum Communication*, pages 151–195. Springer Berlin Heidelberg, Berlin, Heidelberg, 2001.
- [200] R. Horodecki, P. Horodecki, M. Horodecki, and K. Horodecki. Quantum entanglement. *Rev. Modern Phys.*, 81(2):865–942, 2009.
- [201] T. D. Howell. Global properties of tensor rank. *Linear Algebra Appl.*, 22(0):9 – 23, 1978.
- [202] R. Hübener, M. Kleinmann, T.-C. Wei, C. González-Guillén, and O. Gühne. Geometric measure of entanglement for symmetric states. *Phys. Rev. A (3)*, 80(3):032324, 5, 2009.
- [203] I. Ibraghimov. Application of the three-way decomposition for matrix compression. *Numer. Linear Algebra Appl.*, 9(6-7):551–565, 2002.
- [204] L. M. Ioannou, B. C. Travaglione, D. Cheung, and A. K. Ekert. Improved algorithm for quantum separability and entanglement detection. *Phys. Rev. A*, 70:060303, Dec 2004.
- [205] S. Ishizaka. The reduction of the closest disentangled states. *J. Phys. A*, 35(38):8075–8081, 2002.
- [206] M. Ishteva, P.-A. Absil, and P. Van Dooren. Jacobi algorithm for the best low multilinear rank approximation of symmetric tensors. *SIAM J. Matrix Anal. Appl.*, 34(2):651–672, 2013.
- [207] M. Ishteva, P.-A. Absil, S. Van Huffel, and L. De Lathauwer. Best low multilinear rank approximation of higher-order tensors, based on the riemannian trust-region scheme. *SIAM J. Matrix Anal. Appl.*, 32(1):115–135, Feb. 2011.
- [208] M. Ishteva, L. De Lathauwer, P.-A. Absil, and S. Van Huffel. Dimensionality reduction for higher-order tensors: algorithms and applications. *Int. J. Pure Appl. Math.*, 42(3):337–343, 2008.
- [209] M. Ishteva, K. Usevich, and I. Markovsky. Factorization approach to structured low-rank approximation with applications. *SIAM J. Matrix Anal. Appl.*, 35(3):1180–1204, 2014.
- [210] E. T. Jaynes. *Probability theory*. Cambridge University Press, Cambridge, 2003. The logic of science, Edited and with a foreword by G. Larry Bretthorst.
- [211] J.-h. Jiang, H.-l. Wu, Y. Li, and R.-q. Yu. Three-way data resolution by alternating slice-wise diagonalization (asd) method. *Journal of Chemometrics*, 14(1):15–36, 2000.

- [212] N. Jiang, B. Dong, and M. T. Chu. Global rank-1 approximation for order-3 tensors. Tech. report, submitted, North Carolina State University, 2018.
- [213] T. Jiang and N. Sidiropoulos. Kruskal’s permutation lemma and the identification of candecomp/parafac and bilinear models with constant modulus constraints. *Signal Processing, IEEE Transactions on*, 52(9):2625 – 2636, sept. 2004.
- [214] Y.-L. Jiang and X. Kong. On the uniqueness and perturbation to the best rank-one approximation of a tensor. *SIAM J. Matrix Anal. Appl.*, 36(2):775–792, 2015.
- [215] N. Johnston. Norm duality and the cross norm criteria for quantum entanglement. *Linear Multilinear Algebra*, 62(5):648–658, 2014.
- [216] N. Johnston. QETLAB: A MATLAB toolbox for quantum entanglement, version 0.9, Jan. 2016.
- [217] N. Johnston and D. W. Kribs. A family of norms with applications in quantum information theory. *J. Math. Phys.*, 51(8):082202, 16, 2010.
- [218] N. Johnston and D. W. Kribs. Duality of entanglement norms. *Houston J. Math.*, 41(3):831–847, 2015.
- [219] H. Kaiser. The varimax criterion for analytic rotation in factor analysis. *Psychometrika*, 23:187–200, 1958. 10.1007/BF02289233.
- [220] J. Kamm and J. G. Nagy. Optimal Kronecker product approximation of block Toeplitz matrices. *SIAM J. Matrix Anal. Appl.*, 22(1):155–172, 2000.
- [221] A. Kapteyn, H. Neudecker, and T. Wansbeek. An approach to n -mode components analysis. *Psychometrika*, 51(2):269–275, 1986.
- [222] H. B. Keller. Geometrically isolated nonisolated solutions and their approximation. *SIAM J. Numer. Anal.*, 18(5):822–838, 1981.
- [223] J. B. Keller. Closest unitary, orthogonal and Hermitian operators to a given operator. *Math. Mag.*, 48(4):192–197, 1975.
- [224] B. N. Khoromskij. Structured rank- (R_1, \dots, R_D) decomposition of function-related tensors in \mathbb{R}^D . *Comput. Methods Appl. Math.*, 6(2):194–220 (electronic), 2006.
- [225] B. N. Khoromskij and V. Khoromskaia. Low rank Tucker-type tensor approximation to classical potentials. *Cent. Eur. J. Math.*, 5(3):523–550, 2007.
- [226] B. N. Khoromskij, V. Khoromskaia, S. R. Chinnamsetty, and H.-J. Flad. Tensor decomposition in electronic structure calculations on 3D Cartesian grids. *J. Comput. Phys.*, 228(16):5749–5762, 2009.
- [227] H. Kiers. An alternating least squares algorithm for fitting the two- and three-way dedicom model and the idioscal model. *Psychometrika*, 54(3):515–521, 1989.

- [228] H. Kiers. Weighted least squares fitting using ordinary least squares algorithms. *Psychometrika*, 62:251–266, 1997. 10.1007/BF02295279.
- [229] H. A. Kiers and R. A. Harshman. Relating two proposed methods for speedup of algorithms for fitting two- and three-way principal component and related multilinear models. *Chemometrics and Intelligent Laboratory Systems*, 36(1):31 – 40, 1997.
- [230] H. A. L. Kiers. An alternating least squares algorithms for parafac2 and three-way dedicom. *Comput. Stat. Data Anal.*, 16(1):103–118, June 1993.
- [231] H. A. L. Kiers. Joint orthomax rotation of the core and component matrices resulting from three-mode principal components analysis. *J. Classification*, 15(2):245–263, 1998.
- [232] H. A. L. Kiers. A three@step algorithm for candecomp/parafac analysis of large data sets with multicollinearity. *Journal of Chemometrics*, 12(3):155–171, 1998.
- [233] H. A. L. Kiers. Towards a standardized notation and terminology in multiway analysis. *Journal of Chemometrics*, 14(3):105–122, 2000.
- [234] H. A. L. Kiers and A. der Kinderen. A fast method for choosing the numbers of components in Tucker3 analysis. *British J. Math. Statist. Psych.*, 56(1):119–125, 2003.
- [235] H. A. L. Kiers and Y. Takane. Constrained DEDICOM. *Psychometrika*, 58(2):339–355, 1993.
- [236] H. A. L. Kiers, J. M. F. ten Berge, and R. Rocci. Uniqueness of three-mode factor models with sparse cores: the $3 \times 3 \times 3$ case. *Psychometrika*, 62(3):349–374, 1997.
- [237] H. A. L. Kiers, J. M. F. ten Berge, Y. Takane, and J. de Leeuw. A generalization of Takane’s algorithm for DEDICOM. *Psychometrika*, 55(1):151–158, 1990.
- [238] D. E. Knuth. *Art of Computer Programming, Volume 2: Seminumerical Algorithms (3rd Edition)*. Addison-Wesley Professional, 3 edition, Nov. 1997.
- [239] E. Kofidis and P. A. Regalia.
- [240] E. Kofidis and P. A. Regalia. *Tensor Approximation and Signal Processing Applications*. 2001.
- [241] T. G. Kolda. Orthogonal tensor decompositions. *SIAM J. Matrix Anal. Appl.*, 23(1):243–255 (electronic), 2001.
- [242] T. G. Kolda. A counterexample to the possibility of an extension of the Eckart-Young low-rank approximation theorem for the orthogonal rank tensor decomposition. *SIAM J. Matrix Anal. Appl.*, 24(3):762–767, 2003.
- [243] T. G. Kolda. Multilinear operators for higher-order decompositions. Technical Report SAND2006-2081, Sandia National Laboratories, Albuquerque, NM and Livermore, CA, Apr. 2006.

- [244] T. G. Kolda. Numerical optimization for symmetric tensor decomposition. *Math. Program.*, 151(1, Ser. B):225–248, 2015.
- [245] T. G. Kolda and B. W. Bader. Tensor decompositions and applications. *SIAM Rev.*, 51(3):455–500, 2009.
- [246] T. G. Kolda, B. W. Bader, and J. P. Kenny. Higher-order web link analysis using multilinear algebra. In *Proceedings of the Fifth IEEE International Conference on Data Mining, ICDM '05*, pages 242–249, Washington, DC, USA, 2005. IEEE Computer Society.
- [247] W. P. Krijnen. Convergence of the sequence of parameters generated by alternating least squares algorithms. *Comput. Stat. Data Anal.*, 51(2):481–489, Nov. 2006.
- [248] W. P. Krijnen, T. K. Dijkstra, and A. Stegeman. On the non-existence of optimal solutions and the occurrence of “degeneracy” in the CANDECOMP/PARAFAC model. *Psychometrika*, 73(3):431–439, 2008.
- [249] P. Kroonenberg. *Three-mode Principal Component Analysis: Theory and Applications*. M & T series. DSWO Press, 1983.
- [250] P. Kroonenberg. *Applied Multiway Data Analysis*. Wiley Series in Probability and Statistics. Wiley-Interscience, 2008.
- [251] P. Kroonenberg and J. Leeuw. Principal component analysis of three-mode data by means of alternating least squares algorithms. *Psychometrika*, 45:69–97, 1980.
- [252] J. B. Kruskal. Three-way arrays: rank and uniqueness of trilinear decompositions, with application to arithmetic complexity and statistics. *Linear Algebra Appl.*, 18(2):95 – 138, 1977.
- [253] J. B. Kruskal. Multiway data analysis. chapter Rank, decomposition, and uniqueness for 3-way and n-way arrays, pages 7–18. North-Holland Publishing Co., Amsterdam, The Netherlands, The Netherlands, 1989.
- [254] J. B. Kruskal, R. A. Harshman, and M. E. Lundy. Multiway data analysis. chapter How 3-MFA data can cause degenerate parafac solutions, among other relationships, pages 115–122. North-Holland Publishing Co., Amsterdam, The Netherlands, The Netherlands, 1989.
- [255] M. Kuś and K. Życzkowski. Geometry of entangled states. *Phys. Rev. A*, 63:032307, Feb 2001.
- [256] S.-H. Kye. Necessary conditions for optimality of decomposable entanglement witnesses. *Rep. Math. Phys.*, 69(3):419–426, 2012.
- [257] J. D. Laderman. A noncommutative algorithm for multiplying 3×3 matrices using 23 multiplications. *Bull. Amer. Math. Soc.*, 82(1):126–128, 1976.
- [258] W. Landry. Implementing a high performance tensor library. *Sci. Program.*, 11(4):273–290, Dec. 2003.

- [259] J. M. Landsberg. The border rank of the multiplication of 2×2 matrices is seven. *J. Amer. Math. Soc.*, 19(2):447–459, 2006.
- [260] J. M. Landsberg. Geometry and the complexity of matrix multiplication. *Bull. Amer. Math. Soc. (N.S.)*, 45(2):247–284, 2008.
- [261] K. Landsman. Quantum theory and functional analysis, 2019.
- [262] A. N. Langville and W. J. Stewart. The Kronecker product and stochastic automata networks. *J. Comput. Appl. Math.*, 167(2):429–447, 2004.
- [263] A. N. Langville and W. J. Stewart. A kronecker product approximate preconditioner for SANs. *Numer. Linear Algebra Appl.*, 11(8-9):723–752, 2004.
- [264] C. Lawson and R. Hanson. *Solving Least Squares Problems*. Classics in Applied Mathematics. SIAM, 1974.
- [265] C. L. Lawson and R. J. Hanson. *Solving least squares problems*, volume 15 of *Classics in Applied Mathematics*. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 1995.
- [266] D. D. Lee and H. S. Seung. Learning the parts of objects by non-negative matrix factorization. *Nature*, 401(6755):788–791, Oct. 1999.
- [267] D. Leibovici and R. Sabatier. A singular value decomposition of a k-way array for a principal component analysis of multiway data, pta-k. *Linear Algebra Appl.*, 269(1&C3):307 – 329, 1998.
- [268] S. Leurgans and R. T. Ross. Multilinear models: Applications in spectroscopy. *Statistical Science*, 7(3):pp. 289–310, 1992.
- [269] S. Leurgans and R. T. Ross. Multilinear models: applications in spectroscopy. *Statist. Sci.*, 7(3):289–319, 1992. With comments by Jan de Leeuw, Pieter M. Kroonenberg and Donald S. Burdick and a rejoinder by the authors.
- [270] J. Levin. *Three-mode Factor Analysis*. Project on techniques for investigation of structure of individual differences in psychological phenomena. University of Illinois at Urbana-Champaign, 1963.
- [271] C.-G. Li and R. Vidal. A structured sparse plus structured low-rank framework for subspace clustering and completion. *IEEE Trans. Signal Process.*, 64(24):6557–6570, 2016.
- [272] X. Li and X. Fang. Constructing k -Schmidt witnesses for infinite-dimensional systems. *Linear Multilinear Algebra*, 63(4):754–764, 2015.
- [273] L.-H. Lim. Singular values and eigenvalues of tensors: a variational approach. In *Computational Advances in Multi-Sensor Adaptive Processing, 2005 1st IEEE International Workshop on*, pages 129 –132, dec. 2005.
- [274] L.-H. Lim. *Foundations of numerical multilinear algebra: Decomposition and approximation of tensors*. PhD thesis, Stanford University, 2007.

- [275] C.-Y. Lin, Y.-C. Chung, and J.-S. Liu. Efficient data compression methods for multi-dimensional sparse array operations based on the ekmr scheme. *IEEE Trans. Comput.*, 52(12):1640–1646, Dec. 2003.
- [276] C.-Y. Lin, J.-S. Liu, and Y.-C. Chung. Efficient representation scheme for multidimensional array operations. *IEEE Trans. Comput.*, 51(3):327–345, Mar. 2002.
- [277] M. M. Lin, B. Dong, and M. T. Chu. Semi-definite programming techniques for structured quadratic inverse eigenvalue problems. *Numer. Algorithms*, 53(4):419–437, 2010.
- [278] J. Liu, X. Liu, and X. Ma. First-order perturbation analysis of singular vectors in singular value decomposition. *IEEE Transactions on Signal Processing*, 56(7):3044–3049, July 2008.
- [279] N. Liu, B. Zhang, J. Yan, Z. Chen, W. Liu, F. Bai, and L. Chien. Text representation: From vector to tensor. In *Proceedings of the Fifth IEEE International Conference on Data Mining*, ICDM '05, pages 725–728, Washington, DC, USA, 2005. IEEE Computer Society.
- [280] X. Liu and N. Sidiropoulos. Cramer-rao lower bounds for low-rank decomposition of multidimensional arrays. *Trans. Sig. Proc.*, 49(9):2074–2086, Sept. 2001.
- [281] S. Łojasiewicz. Une propriété topologique des sous-ensembles analytiques réels. In *Les Équations aux Dérivées Partielles (Paris, 1962)*, pages 87–89. Éditions du Centre National de la Recherche Scientifique, Paris, 1963.
- [282] S. Łojasiewicz and M.-A. Zurro. On the gradient inequality. *Bull. Polish Acad. Sci. Math.*, 47(2):143–145, 1999.
- [283] D. G. Luenberger. *Optimization by vector space methods*. John Wiley & Sons, Inc., New York-London-Sydney, 1969.
- [284] M. E. Lundy, R. E. Harshamn, and J. B. Kruskal. Multiway data analysis. chapter A two-stage procedure incorporating good features of both trilinear and quadrilinear models, pages 123–130. North-Holland Publishing Co., Amsterdam, The Netherlands, The Netherlands, 1989.
- [285] M. E. Lundy, R. A. Harshman, P. Paatero, and L. C. Swartzman. Application of the 3-way dedicom model to skew-symmetric data for paired preference ratings of treatments for chronic back pain. Technical report, 2003.
- [286] J. Malick. A dual approach to semidefinite least-squares problems. *SIAM J. Matrix Anal. Appl.*, 26(1):272–284, 2004.
- [287] G. I. Marchuk. Construction of adjoint operators in non-linear problems of mathematical physics. *Sbornik: Mathematics*, 189(10):1505, 1998.
- [288] C. D. M. Martin and C. F. Van Loan. A jacobi-type method for computing orthogonal tensor decompositions. *SIAM J. Matrix Anal. Appl.*, 30(3):1219–1232, Sept. 2008.
- [289] J. M. Martínez. Local convergence theory of inexact Newton methods based on structured least change updates. *Math. Comp.*, 55(191):143–167, 1990.

- [290] MathWorks. *MATLAB Optimization Toolbox, Version 8.1*. Natick, MA, USA, 2018.
- [291] M. Melucci. *Introduction to information retrieval and quantum mechanics*, volume 35 of *The Information Retrieval Series*. Springer, Heidelberg, 2015.
- [292] B. Merlet and M. Pierre. Convergence to equilibrium for the backward Euler scheme and applications. *Commun. Pure Appl. Anal.*, 9(3):685–702, 2010.
- [293] F. Mintert, A. R. R. Carvalho, M. Kuś, and A. Buchleitner. Measures and dynamics of entangled states. *Phys. Rep.*, 415(4):207–259, 2005.
- [294] A. Miranowicz and S. Ishizaka. Closed formula for the relative entropy of entanglement. *Phys. Rev. A (3)*, 78(3):032310, 7, 2008.
- [295] B. C. Mitchell and D. S. Burdick. Slowly converging parafac sequences: Swamps and two-factor degeneracies. *Journal of Chemometrics*, 8(2):155–168, 1994.
- [296] F. Miwakeichi, E. Martínez-Montes, P. A. Valdés-Sosa, N. Nishiyama, H. Mizuhara, and Y. Yamaguchi. Decomposing EEG data into space-time-frequency components using parallel factor analysis. *NeuroImage*, 22(3):1035 – 1045, 2004.
- [297] F. Miwakeichi, P. A. Valdes-Sosa, E. Aubert-Vazquez, J. B. Bayard, J. Watanabe, H. Mizuhara, and Y. Yamaguchi. Neural information processing. chapter Decomposing EEG Data into Space-Time-Frequency Components Using Parallel Factor Analysis and Its Relation with Cerebral Blood Flow, pages 802–810. Springer-Verlag, Berlin, Heidelberg, 2008.
- [298] J. Mocks. Topographic components model for event-related potentials and some biophysical considerations. *Biomedical Engineering, IEEE Transactions on*, 35(6):482–484, June 1988.
- [299] M. J. Mohlenkamp. Musings on multilinear fitting. *Linear Algebra Appl.*, 438(2):834–852, 2013.
- [300] J. J. Moré and D. C. Sorensen. Computing a trust region step. *SIAM J. Sci. Statist. Comput.*, 4(3):553–572, 1983.
- [301] A. P. Morgan, A. J. Sommese, and L. T. Watson. Finding all isolated solutions to polynomial systems using HOMPACk. *ACM Trans. Math. Software*, 15(2):93–122, 1989.
- [302] M. Mørup, L. K. Hansen, and S. M. Arnfred. Algorithms for sparse nonnegative Tucker decompositions. *Neural Comput.*, 20(8):2112–2131, Aug. 2008.
- [303] T. Murakami, J. Ten Berge, and H. Kiers. A case of extreme simplicity of the core matrix in three-mode principal components analysis. *Psychometrika*, 63:255–261, 1998. 10.1007/BF02294854.
- [304] D. Muti and S. Bourennane. Multidimensional filtering based on a tensor approach. *Signal Process.*, 85(12):2338–2353, Dec. 2005.

- [305] J. Nagy and M. Kilmer. Kronecker product approximation for preconditioning in three-dimensional imaging applications. *Image Processing, IEEE Transactions on*, 15(3):604–613, march 2006.
- [306] M. Nakahara and T. Ohmi. *Quantum computing*. CRC Press, Boca Raton, FL, 2008. From linear algebra to physical realizations.
- [307] M. Narasimhan. *Principles of continuum mechanics*. Wiley-Interscience publication. Wiley, 1993.
- [308] C. Navasca, L. De Lathauwer, and S. Kindermann. Swamp reducing technique for tensor decomposition. In *Proc. 16th European Signal Processing Conference (EUSIPCO 2008)*, Lausanne, Switzerland, Aug. 25–29 2008.
- [309] G. Ni, L. Qi, and M. Bai. Geometric measure of entanglement and U-eigenvalues of tensors. *SIAM J. Matrix Anal. Appl.*, 35(1):73–87, 2014.
- [310] G. Ni and Y. Wang. On the best rank-1 approximation to higher-order symmetric tensors. *Math. Comput. Modelling*, 46(9-10):1345–1352, 2007.
- [311] M. A. Nielsen and I. L. Chuang. *Quantum Computation and Quantum Information: 10th Anniversary Edition*. Cambridge University Press, 2010.
- [312] D. Nion and L. De Lathauwer. Fast communication: An enhanced line search scheme for complex-valued tensor decompositions. application in ds-cdma. *Signal Process.*, 88(3):749–755, Mar. 2008.
- [313] J. M. Ortega. *Numerical analysis*, volume 3 of *Classics in Applied Mathematics*. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, second edition, 1990. A second course.
- [314] P. Paatero. A weighted non-negative least squares algorithm for three-way PARAFAC factor analysis. *Chemometrics and Intelligent Laboratory Systems*, 38(2):223 – 242, 1997.
- [315] P. Paatero. The multilinear engine: A table-driven, least squares program for solving multilinear problems, including the n-way parallel factor analysis model. *Journal of Computational and Graphical Statistics*, 8(4):pp. 854–888, 1999.
- [316] P. Paatero. Construction and analysis of degenerate parafac models. *Journal of Chemometrics*, 14(3):285–299, 2000.
- [317] P. Paatero and U. Tapper. Positive matrix factorization: A non-negative factor model with optimal utilization of error estimates of data values. *Environmetrics*, 5(2):111–126, 1994.
- [318] R. S. Palais. The principle of symmetric criticality. *Comm. Math. Phys.*, 69(1):19–30, 1979.
- [319] A. Peres. Separability criterion for density matrices. *Phys. Rev. Lett.*, 77(8):1413–1415, 1996.
- [320] V. Pereyra and G. Scherer. Efficient computer manipulation of tensor products with applications to multidimensional approximation. *Math. Comput.*, 27:595–605, 1973.

- [321] M. Pierre. Quelques applications de l'inégalité de Lojasiewicz à des discrétisations d'EDP. SMAI, 2011.
- [322] M. B. Płbnió and S. Virmani. An introduction to entanglement measures. *Quantum Info. Comput.*, 7(1):1–51, Jan. 2007.
- [323] L. Qi. Some simple estimates for singular values of a matrix. *Linear Algebra Appl.*, 56(0):105 – 119, 1984.
- [324] L. Qi. Eigenvalues of a real supersymmetric tensor. *J. Symb. Comput.*, 40(6):1302–1324, Dec. 2005.
- [325] L. Qi. Eigenvalues and invariants of tensors. *Journal of Mathematical Analysis and Applications*, 325:1363–1377, 2007.
- [326] L. Qi. The best rank-one approximation ratio of a tensor space. *SIAM J. Matrix Analysis Applications*, pages 430–442, 2011.
- [327] L. Qi. The best rank-one approximation ratio of a tensor space. *SIAM J. Matrix Anal. Appl.*, 32(2):430–442, 2011.
- [328] L. Qi, W. Sun, and Y. Wang. Numerical multilinear algebra and its applications. *Front. Math. China*, 2:501 – 526, 2007.
- [329] B. R. and A. C.A. Improving the speed of multiway algorithms - part ii: Compression. *Chemometrics and Intelligent Laboratory Systems*, 42(1):105–113, 1998.
- [330] M. Rajih, P. Comon, and R. A. Harshman. Enhanced line search: A novel method to accelerate parafac. *SIAM J. Matrix Anal. Appl.*, 30(3):1128–1147, Sept. 2008.
- [331] R. Raussendorf and H. J. Briegel. A one-way quantum computer. *Phys. Rev. Lett.*, 86:5188–5191, May 2001.
- [332] W. S. Rayens and B. C. Mitchell. Two-factor degeneracies and a stabilization of parafac. *Chemometrics and Intelligent Laboratory Systems*, 38(2):173 – 181, 1997.
- [333] P. Regalia and E. Kofidis. The higher-order power method revisited: convergence proofs and effective initialization. In *Acoustics, Speech, and Signal Processing, 2000. ICASSP '00. Proceedings. 2000 IEEE International Conference on*, volume 5, pages 2709 –2712 vol.5, 2000.
- [334] P. A. Regalia. Monotonically convergent algorithms for symmetric tensor approximation. *Linear Algebra Appl.*, 438(2):875–890, 2013.
- [335] M. J. Reynolds, A. Doostan, and G. Beylkin. Randomized alternating least squares for canonical tensor decompositions: application to a PDE with random data. *SIAM J. Sci. Comput.*, 38(5):A2634–A2664, 2016.
- [336] R. Rocci. A general algorithm to fit constrained dedicom models. *Statistical Methods and Applications*, 13:139–150, 2004.

- [337] T. Rohwedder and A. Uschmajew. On local convergence of alternating schemes for optimization of convex problems in the tensor train format. *SIAM J. Numer. Anal.*, 51(2):1134–1162, 2013.
- [338] A. Ruhe. Algorithms for the nonlinear eigenvalue problem. *SIAM J. Numer. Anal.*, 10:674–689, 1973.
- [339] A. Rutkowski and P. Horodecki. Tensor product extension of entanglement witnesses and their connection with measurement-device-independent entanglement witnesses. *Physics Letters A*, 378(30):2043 – 2047, 2014.
- [340] E. Sanchez and B. R. Kowalski. Tensorial resolution: A direct trilinear decomposition. *Journal of Chemometrics*, 4(1):29–45, 1990.
- [341] B. Savas. Analyses and tests of handwritten digit recognition algorithms, 2003.
- [342] B. Savas and L. Eldén. Handwritten digit classification using higher order singular value decomposition. *Pattern Recognition*, 40(3):993 – 1003, 2007.
- [343] B. Savas and L.-H. Lim. Quasi-Newton methods on Grassmannians and multilinear approximations of tensors. *SIAM J. Sci. Comput.*, 32(6):3352–3393, 2010.
- [344] A. Schönhage. Partial and total matrix multiplication. *SIAM Journal on Computing*, 10(3):434–455, 1981.
- [345] E. Schost and P.-J. Spaenlehauer. A quadratically convergent algorithm for structured low-rank approximation. *Found. Comput. Math.*, 16(2):457–492, 2016.
- [346] A. Shashua and T. Hazan. Non-negative tensor factorization with applications to statistics and computer vision. In *Proceedings of the 22nd international conference on Machine learning*, ICML '05, pages 792–799, New York, NY, USA, 2005. ACM.
- [347] A. Shashua and A. Levin. Linear image coding for regression and classification using the tensor-rank principle. In *Proceedings of the 2001 IEEE Computer Society Conference on Computer Vision and Pattern Recognition, 2001*, volume 1, pages I42–49, 2001.
- [348] A. Shashua, R. Zass, and T. Hazan. Multi-way clustering using super-symmetric non-negative tensor factorization. In A. Leonardis, H. Bischof, and A. Pinz, editors, *ECCV (4)*, volume 3954 of *Lecture Notes in Computer Science*, pages 595–608. Springer, 2006.
- [349] P. W. Shor. Polynomial-time algorithms for prime factorization and discrete logarithms on a quantum computer. *SIAM J. Comput.*, 26(5):1484–1509, 1997.
- [350] P. W. Shor. Polynomial-time algorithms for prime factorization and discrete logarithms on a quantum computer. *SIAM Rev.*, 41(2):303–332, 1999.
- [351] M. Shub and S. Smale. Complexity of Bezout’s theorem. II. Volumes and probabilities. In *Computational algebraic geometry (Nice, 1992)*, volume 109 of *Progr. Math.*, pages 267–285. Birkhäuser Boston, Boston, MA, 1993.

- [352] N. Sidiropoulos, R. Bro, and G. Giannakis. Parallel factor analysis in sensor array processing. *Signal Processing, IEEE Transactions on*, 48(8):2377–2388, aug 2000.
- [353] N. Sidiropoulos and R. Budampati. Khatri-rao space-time codes. *Signal Processing, IEEE Transactions on*, 50(10):2396–2407, oct 2002.
- [354] N. Sidiropoulos, G. Giannakis, and R. Bro. Blind parafac receivers for ds-cdma systems. *Signal Processing, IEEE Transactions on*, 48(3):810–823, mar 2000.
- [355] N. D. Sidiropoulos and R. Bro. On the uniqueness of multilinear decomposition of n-way arrays. *Journal of Chemometrics*, 14(3):229–239, 2000.
- [356] A. I. Singh. Role of partial transpose and generalized Choi maps in quantum dynamical semigroups involving separable and entangled states. *Electron. J. Linear Algebra*, 29:156–193, 2015.
- [357] A. Smilde, R. Bro, and P. Geladi. *Multi-way Analysis with Applications in the Chemical Sciences*. J. Wiley, 2004.
- [358] A. Smilde, R. Bro, and P. Geladi. *Multi-way Analysis: Applications in the Chemical Sciences*. Wiley InterScience online books. Wiley, 2005.
- [359] A. K. Smilde, Y. Wang, and B. R. Kowalski. Theory of medium-rank second-order calibration with restricted-tucker models. *Journal of Chemometrics*, 8(1):21–36, 1994.
- [360] A. J. Sommese and C. W. Wampler, II. *The numerical solution of systems of polynomials arising in engineering and science*. World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2005.
- [361] L. Sorber, M. Van Barel, and L. De Lathauwer. Unconstrained optimization of real functions in complex variables. *SIAM J. Optim.*, 22(3):879–898, 2012.
- [362] L. Sorber, M. Van Barel, and L. De Lathauwer. Optimization-based algorithms for tensor decompositions: canonical polyadic decomposition, decomposition in rank- $(L_r, L_r, 1)$ terms, and a new generalization. *SIAM J. Optim.*, 23(2):695–720, 2013.
- [363] M. Sørensen, L. D. Lathauwer, P. Comon, S. Icart, and L. Deneire. Canonical Polyadic Decomposition with a Columnwise Orthonormal Factor Matrix. *SIAM J. Matrix Anal. Appl.*, 33(4):1190–1213, 2012.
- [364] A. Stegeman. Degeneracy in candecomp/parafac explained for $p \times p \times 2$ arrays of rank $p + 1$ or higher. *Psychometrika*, 71:483–501, 2006.
- [365] A. Stegeman. Degeneracy in candecomp/parafac and indscal explained for several three-sliced arrays with a two-valued typical rank. *Psychometrika*, 72:601–619, 2007.
- [366] A. Stegeman. Low-rank approximation of generic $p \times q \times 2$ arrays and diverging components in the Candecomp/Parafac model. *SIAM J. Matrix Anal. Appl.*, 30(3):988–1007, 2008.

- [367] A. Stegeman, J. Berge, and L. Lathauwer. Sufficient conditions for uniqueness in candecomp/parafac and indscal with random component matrices. *Psychometrika*, 71:219–229, 2006.
- [368] A. Stegeman and P. Comon. Subtracting a best rank-1 approximation may increase tensor rank. *Linear Algebra Appl.*, 433(7):1276–1300, 2010.
- [369] A. Stegeman and N. D. Sidiropoulos. On kruskal’s uniqueness condition for the candecomp/parafac decomposition. *Linear Algebra Appl.*, 420(2&C3):540 – 552, 2007.
- [370] V. Strassen. Gaussian elimination is not optimal. *Numerische Mathematik*, 13:354–356, 1969. 10.1007/BF02165411.
- [371] J. Sun, S. Papadimitriou, and P. S. Yu. Window-based tensor analysis on high-dimensional and multi-aspect streams. In *Proceedings of the Sixth International Conference on Data Mining*, ICDM ’06, pages 1076–1080, Washington, DC, USA, 2006. IEEE Computer Society.
- [372] J. Sun, D. Tao, and C. Faloutsos. Beyond streams and graphs: dynamic tensor analysis. In *Proceedings of the 12th ACM SIGKDD international conference on Knowledge discovery and data mining*, KDD ’06, pages 374–383, New York, NY, USA, 2006. ACM.
- [373] J.-T. Sun, H.-J. Zeng, H. Liu, Y. Lu, and Z. Chen. Cubesvd: a novel approach to personalized web search. In *Proceedings of the 14th international conference on World Wide Web*, WWW ’05, pages 382–390, New York, NY, USA, 2005. ACM.
- [374] A. Takayama. *Mathematical economics*. Cambridge University Press, Cambridge, second edition, 1985.
- [375] S. Tamaryan. Completely mixed state is a critical point for three-qubit entanglement. *Phys. Lett. A*, 375(23):2224–2229, 2011.
- [376] J. ten Berge. Kruskal’s polynomial for $2 \times 2 \times 2$ arrays and a generalization to $2 \times 2 \times n$ arrays. *Psychometrika*, 56:631–636, 1991. 10.1007/BF02294495.
- [377] J. ten Berge. The typical rank of tall three-way arrays. *Psychometrika*, 65:525–532, 2000. 10.1007/BF02296342.
- [378] J. ten Berge and H. Kiers. Some uniqueness results for PARAFAC2. *Psychometrika*, 61:123–132, 1996. 10.1007/BF02296962.
- [379] J. ten Berge and N. Sidiropoulos. On uniqueness in candecomp/parafac. *Psychometrika*, 67:399–409, 2002. 10.1007/BF02294992.
- [380] J. M. ten Berge, N. D. Sidiropoulos, and R. Rocci. Typical rank and indscal dimensionality for symmetric three-way arrays of order $i \times 2 \times 2$ or $i \times 3 \times 3$. *Linear Algebra Appl.*, 388(0):363 – 377, 2004. Tenth Special Issue (Part 1) on Linear Algebra and Statistics.
- [381] J. M. F. ten Berge. *Least squares optimization in multivariate analysis*, volume 25 of *M&T Series*. D.S.W.O. Press, Leiden, 1993.

- [382] J. M. F. ten Berge. Partial uniqueness in CANDECOMP/PARAFAC. *Journal of Chemometrics*, 18(1):12–16, 2004.
- [383] J. M. F. ten Berge, H. A. L. Kiers, and J. de Leeuw. Explicit CANDECOMP/PARAFAC solutions for a contrived $2 \times 2 \times 2$ array of rank three. *Psychometrika*, 53(4):579–583, 1988.
- [384] B. M. Terhal. Detecting quantum entanglement. *Theoret. Comput. Sci.*, 287(1):313–335, 2002. Natural computing.
- [385] B. M. Terhal and P. Horodecki. Schmidt number for density matrices. *Phys. Rev. A (3)*, 61(4):040301, 4, 2000.
- [386] W. Thirring, R. A. Bertlmann, P. Köhler, and H. Narnhofer. Entanglement or separability: the choice of how to factorize the algebra of a density matrix. *The European Physical Journal D*, 64(2):181–196, Oct 2011.
- [387] M. J. Todd. Semidefinite optimization. *Acta Numer.*, 10:515–560, 2001.
- [388] G. Tomasi. *Practical and Computational Aspects in Chemometric Data Analysis: Ph.D. Dissertation*. Department of Food Science, Royal Veterinary and Agricultural University, 2006.
- [389] G. Tomasi and R. Bro. PARAFAC and missing values. *Chemometrics and Intelligent Laboratory Systems*, 75(2):163 – 180, 2005.
- [390] G. Tomasi and R. Bro. A comparison of algorithms for fitting the PARAFAC model. *Comput. Stat. Data Anal.*, 50(7):1700–1734, Apr. 2006.
- [391] L. Tucker. Some mathematical notes on three-mode factor analysis. *Psychometrika*, 31:279–311, 1966.
- [392] L. R. Tucker. Implications of factor analysis of three-way matrices for measurement of change. In C. W. Harris, editor, *Problems in measuring change.*, pages 122–137. University of Wisconsin Press, Madison WI, 1963.
- [393] L. R. Tucker. The extension of factor analysis to three-dimensional matrices. In H. Gulliksen and N. Frederiksen, editors, *Contributions to mathematical psychology.*, pages 110–127. Holt, Rinehart and Winston, New York, 1964.
- [394] Z. Ugray, L. Lasdon, J. Plummer, F. Glover, J. Kelly, and R. Martí. Scatter search and local NLP solvers: a multistart framework for global optimization. *INFORMS J. Comput.*, 19(3):328–340, 2007.
- [395] A. Uschmajew. Local convergence of the alternating least squares algorithm for canonical tensor approximation. *SIAM J. Matrix Anal. Appl.*, 33(2):639–652, 2012.
- [396] A. Uschmajew. A new convergence proof for the higher-order power method and generalizations. *Pac. J. Optim.*, 11(2):309–321, 2015.
- [397] C. F. Van Loan. The ubiquitous Kronecker product. *J. Comput. Appl. Math.*, 123(1-2):85–100, 2000.

- [398] C. F. Van Loan. Structured matrix problems from tensors. In *Exploiting Hidden Structure in Matrix Computations: Algorithms and Applications*, volume 2173 of *Lecture Notes in Math.*, pages 1–63. Springer, Cham, 2016.
- [399] C. F. Van Loan and N. Pitsianis. Approximation with Kronecker products. In *Linear algebra for large scale and real-time applications (Leuven, 1992)*, volume 232 of *NATO Adv. Sci. Inst. Ser. E Appl. Sci.*, pages 293–314. Kluwer Acad. Publ., Dordrecht, 1993.
- [400] L. Vandenberghe and S. Boyd. Semidefinite programming. *SIAM Rev.*, 38(1):49–95, 1996.
- [401] N. Vannieuwenhoven, R. Vandebril, and K. Meerbergen. A new truncation strategy for the higher-order singular value decomposition. *SIAM J. Sci. Comput.*, 34(2):A1027–A1052, 2012.
- [402] M. Vasilescu. Human motion signatures: analysis, synthesis, recognition. In *Pattern Recognition, 2002. Proceedings. 16th International Conference on*, volume 3, pages 456 – 460 vol.3, 2002.
- [403] M. Vasilescu and D. Terzopoulos. Multilinear image analysis for facial recognition. In *Pattern Recognition, 2002. Proceedings. 16th International Conference on*, volume 2, pages 511 – 514 vol.2, 2002.
- [404] M. A. O. Vasilescu and D. Terzopoulos. Multilinear subspace analysis of image ensembles. In *CVPR (2)'03*, pages 93–99, 2003.
- [405] M. A. O. Vasilescu and D. Terzopoulos. Tensortextures: multilinear image-based rendering. *ACM Trans. Graph.*, 23(3):336–342, Aug. 2004.
- [406] M. A. O. Vasilescu and D. Terzopoulos. Multilinear independent components analysis. In *Proceedings of the 2005 IEEE Computer Society Conference on Computer Vision and Pattern Recognition (CVPR'05) - Volume 1 - Volume 01*, CVPR '05, pages 547–553, Washington, DC, USA, 2005. IEEE Computer Society.
- [407] L. Vega-Montoto and P. D. Wentzell. Maximum likelihood parallel factor analysis (ML-PARAFAC). *Journal of Chemometrics*, 17(4):237–253, 2003.
- [408] N. Vervliet, O. Debals, L. Sorber, M. Van Barel, and L. De Lathauwer. Tensorlab 3.0, Mar. 2016. Available online.
- [409] D. Vlastic, M. Brand, H. Pfister, and J. Popović. Face transfer with multilinear models. *ACM Trans. Graph.*, 24(3):426–433, July 2005.
- [410] D. Vlastic, M. Brand, H. Pfister, and J. Popovic. Face transfer with multilinear models. In *ACM SIGGRAPH 2006 Courses*, SIGGRAPH '06, New York, NY, USA, 2006. ACM.
- [411] M. D. Vos, A. Vergult, L. D. Lathauwer, W. D. Clercq, S. V. Huffel, P. Dupont, A. Palmiini, and W. V. Paesschen. Canonical decomposition of ictal scalp eeg reliably detects the seizure onset zone. *NeuroImage*, 37(3):844 – 854, 2007.
- [412] V. Vu. Singular vectors under random perturbation. *Random Structures Algorithms*, 39(4):526–538, 2011.

- [413] E. Wakakuwa, A. Soeda, and M. Murao. A coding theorem for bipartite unitaries in distributed quantum computation. *IEEE Trans. Inform. Theory*, 63(8):5372–5403, 2017.
- [414] H. Wang and N. Ahuja. Facial expression decomposition. In *Proceedings of the Ninth IEEE International Conference on Computer Vision - Volume 2, ICCV '03*, pages 958–, Washington, DC, USA, 2003. IEEE Computer Society.
- [415] H. Wang and N. Ahuja. Compact representation of multidimensional data using tensor rank-one decomposition. In *Proceedings of the Pattern Recognition, 17th International Conference on (ICPR'04) Volume 1 - Volume 01, ICPR '04*, pages 44–47, Washington, DC, USA, 2004. IEEE Computer Society.
- [416] L. Wang and M. T. Chu. On the global convergence of the alternating least squares method for rank-one approximation to generic tensors. *SIAM J. Matrix Anal. Appl.*, 35(3):1058–1072, 2014.
- [417] L. Wang, M. T. Chu, and B. Yu. Orthogonal low rank tensor approximation: alternating least squares method and its global convergence. *SIAM J. Matrix Anal. Appl.*, 36(1):1–19, 2015.
- [418] W. C. Waterhouse. Do symmetric problems have symmetric solutions? *Amer. Math. Monthly*, 90(6):378–387, 1983.
- [419] P.-Å. Wedin. Perturbation bounds in connection with singular value decomposition. *Nordisk Tidskr. Informationsbehandling (BIT)*, 12:99–111, 1972.
- [420] T.-C. Wei and S. Severini. Matrix permanent and quantum entanglement of permutation invariant states. *J. Math. Phys.*, 51(9):092203, 11, 2010.
- [421] M. Welling and M. Weber. Positive tensor factorization. *Pattern Recogn. Lett.*, 22(12):1255–1261, Oct. 2001.
- [422] R. F. Werner. Quantum states with einstein-podolsky-rosen correlations admitting a hidden-variable model. *Phys. Rev. A*, 40:4277–4281, Oct 1989.
- [423] M. M. Wilde. *Quantum information theory*. Cambridge University Press, Cambridge, second edition, 2017.
- [424] B. M. Wise, N. B. Gallagher, and E. B. Martin. Application of PARAFAC2 to fault detection and diagnosis in semiconductor etch. *Journal of Chemometrics*, 15(4):285–298, 2001.
- [425] K. Wright. Differential equations for the analytic singular value decomposition of a matrix. *Numer. Math.*, 3(2):283–295, 1992.
- [426] M. Wschebor. On the Kostlan-Shub-Smale model for random polynomial systems. Variance of the number of roots. *J. Complexity*, 21(6):773–789, 2005.
- [427] H.-L. Wu, M. Shibukawa, and K. Oguma. An alternating trilinear decomposition algorithm with application to calibration of HPLC-CDAD for simultaneous determination of overlapped chlorinated aromatic hydrocarbons. *Journal of Chemometrics*, 12(1):1–26, 1998.

- [428] S.-J. Wu and M. T. Chu. Markov chains with memory, tensor formulation, and the dynamics of power iteration. *Appl. Math. Comput.*, 303:226–239, 2017.
- [429] S.-J. Wu and M. T. Chu. Solving an inverse eigenvalue problem with triple constraints on eigenvalues, singular values, and diagonal elements. *Inverse Problems*, 33(8):085003, 21, 2017.
- [430] B. Xhabli. The super operator system structures and their applications in quantum entanglement theory. *J. Funct. Anal.*, 262(4):1466–1497, 2012.
- [431] Y. Yang, S. Hu, L. De Lathauwer, and J. A. Suykens. Convergence study of block singular value maximization methods for rank-1 approximation to higher order tensors. Technical report, Internal Report 16-149, ESAT-SISTA, KU Leuven, 2016.
- [432] E. Zander and H. G. Matthies. Tensor product methods for stochastic problems. *PAMM*, 7(1):2040067–2040068, 2007.
- [433] E. Zhang, J. Hays, and G. Turk. Interactive tensor field design and visualization on surfaces. *IEEE Transactions on Visualization and Computer Graphics*, 13, 2007.
- [434] L.-H. Zhang and M. T. Chu. Computing absolute maximum correlation. *IMA J. Numer. Anal.*, 32(1):163–184, 2012.
- [435] L.-H. Zhang, L.-Z. Liao, and L.-M. Sun. Towards the global solution of the maximal correlation problem. *J. of Global Optimization*, 49(1):91–107, Jan. 2011.
- [436] T. Zhang and G. H. Golub. Rank-one approximation to high order tensors. *SIAM J. Matrix Anal. Appl.*, 23(2):534–550, Feb. 2001.
- [437] X. Zhang, C. Ling, and L. Qi. The best rank-1 approximation of a symmetric tensor and related spherical optimization problems. *SIAM J. Matrix Anal. Appl.*, 33(3):806–821, 2012.